

Adult Organ Preservation (For Brain Dead Patients Only)

INSTRUCTIONS FOR THE USE OF THIS ORDER:

Physician signature at bottom of form enacts all pre-marked orders. To deselect a pre-marked order draw a line through it. To select an order that is not pre-selected place an "X" or "✓" inside the unmarked box.

- Transfer Care to Gift of Life Michigan
- Discontinue all previous orders and medications
- Notify House Supervisor/Manager of donation
- Notify On Call Specialty Physicians
 - _____ (Cardiology) for Heart Catheterization
 - _____ (Cardiology) for Echocardiogram
 - _____ (Pulmonology) for Bronchoscopy
 - _____ (Intensivist) for Swan (PA line) insertion
 - _____ (Intensivist) for Central Line insertion
 - _____ (Intensivist) for Arterial Line insertion
 - _____ (Intensivist) for Biopsy
 - _____ (Pathology) for Biopsy interpretation
 - _____ (Radiology) for CXR Interpretation
 - _____ (Other) _____

NURSING

- Vital signs every hour
- Urine output every hour
- Maintain body temp 96-98.5 degrees Fahrenheit-use warming or cooling blanket
- Nasogastric tube to low continuous suction
- Sequential compression devices
- Central venous pressure
- Pulmonary artery pressure
- Foley catheter to dependant drainage
- Suction every 1-2 hours
- Turn every 1-2 hours

LABS

- CBC, Electrolytes, BUN, Cr, Random glucose, calcium, magnesium, phosphorus, ionized calcium, GFR, ALT, AST, Total and Direct Bilirubin, alk phos, LDH, Total protein, albumin, PT, PTT, INR, amylase, lipase, troponin, CPK, CPK-MB, GGT x 1 **stat and every 4 hours**
- ABG **stat and every 4 hours**
- U/A
- Sputum gram stain

TESTING

- EKG
- Portable CXR stat R/O Infiltrate
- Echocardiogram
- Bronchoscopy
- Cardiac Catheterization
- Liver Biopsy

RESPIRATORY CARE

- Ventilator Settings:
Mode: _____ Rate: _____ Vt: _____ FiO2: _____ PEEP: _____
- Chest percussions every 1-2 hours by bed

Key:
 \leq = less than or equal to
 \geq = greater than or equal to
 $<$ = less than
 $>$ = greater than

MEDICATIONS

- Maintenance IV fluid _____ at _____ mL/hr
 - Give the following 4 medications in rapid succession at _____ (time) after first brain death exam is complete:
 - Vasopressin (Pitressin) 1 unit IVP x 1
 - Methylprednisolone (Solu-Medrol®) 15 mg/kg IVP x 1
 - Insulin Regular (Novolin) 10 units IVP x 1
 - Levothyroxine (Synthroid,T-4) 20 mcg IVP x 1
 - Vasopressin (Pitressin) 20 units/150 mL D5W
 - 0.5 – 4.0 units/hour IV infusion. Titrate to maintain an SVR 800 - 1200 dyne/sec cm⁵
 - OR**
 - Titrate to urine output 0.5-3.0 cc/kg/hour
 - Levothyroxine (Synthroid,T-4) 200 mcg/ 500 mL 0.9% NaCl at 25mL/hour (10mcg)
 - Methylprednisolone (Solu-Medrol®) 15 mg/kg IVP every 6 hours
 - Dopamine 400 mg/250mL D5W 2mcg/kg/min, titrate by 2.5 mcg/kg/min every 5 min up to 20 mcg/kg/min to SBP >____ or MAP>____
 - Epinephrine (Adrenalin®) 1 mg/250 mL D5W 2 mcg/min, titrate by 1 mcg/min every 5 min up to 10 mcg/min to SBP>____ or MAP>____
 - Norepinephrine (Levophed®) 4 mg/250 mL D5W 2 mcg/min, titrate by 1 mcg/min every 5 min up to 12 mcg/min to SBP>____ or MAP>____
 - Dobutamine (Dobutrex®) 500 mg/250 mL D5W 2 mcg/kg/min, titrate by 2.5 mcg/kg/min every 5 min up to 20 mcg/kg/min to maintain cardiac index > _____
 - Phenylephrine (Neosynephrine®) 10 mg/ 250 mL D5W 40 mcg/min, titrate by 5 mcg/min every 5 min up to 180 mcg/min to SBP > _____ or MAP >_____
 - Albuterol (Proventil®) 2.5 mg unit dose nebulized every 4 hours
 - Ipratropium (Atrovent®) 0.5 mg unit dose nebulized every 4 hours
 - Acetylcysteine (Mucomyst®) 20% 1,000 mg (5mL) via endotracheal tube every 4 hours PRN-secretions
 - Cefazolin (Kefzol®) _____ gm IV every _____ hrs
 - Clindamycin (Cleocin®) _____ mg IV every _____ hrs
 - Heparin 5000 units SQ every 8 hrs
 - Prilosec 40 mg NGT every day
- The following 2 medications may only be given at _____ (time):**
- Naloxone (Narcan®) 8 mg IVP x 1
 - Vecuronium (Norcuron®) 10 mg IVP x 1. Give with Naloxone

For Use As A Guideline Only

Gift of Life Michigan
Signature _____ Date _____ Time _____

Noted _____ Date _____ Time _____